arXiv:1212.2413v2 [math.OA] 12 Dec 2012

AN ERRATUM TO "PROPERTY (RD) FOR HECKE PAIRS"

VAHID SHIRBISHEH

Let (G, H) be a Hecke pair. The popular version of the reduced Hecke C^*-algebra associated to (G, H) is based on the right action of $\mathcal{H}(G, H)$ (by the right convolution) on $\ell^2(G/H)$. In our paper [1], since we wanted the Hecke algebra $\mathcal{H}(G, H)$ acts from the left on a Hilbert space, we used the the left action of $\mathcal{H}(G, H)$ (by the left convolution) on $\ell^2(H\backslash G)$. To this work, we must have used the following formula for the convolution product instead of the one appearing in Definition 1.1 of [1]:

$$f_1 * f_2(g) := \sum_{\gamma \in < H\backslash G >} f_1(g\gamma^{-1}) f_2(\gamma), \tag{0.1}$$

for all $f_1, f_2 \in \mathcal{H}(G, H)$ and $g \in < G//H >$ (or similarly for all $f_1 \in \mathcal{H}(G, H)$, $f_2 \in \ell^2(H\backslash G)$ and $g \in < H\backslash G >$). One notes that these two formulas are equivalent for groups. But in the setting of Hecke pairs, our previous formula for the convolution product does not define a structure of an involutive algebra on $\mathcal{H}(G, H)$.

By applying this change to every occurrence of the convolution product in [1], (precisely, in Pages 174, 182, 183), every statement is still valid.

To show that these two approaches for defining a reduced Hecke C^*-algebra are equivalent, let us denote the left and right actions of $\mathcal{H}(G, H)$ on $\ell^2(H\backslash G)$ and $\ell^2(G/H)$ by λ_l and λ_r, respectively. Define $U : \ell^2(H\backslash G) \to \ell^2(G/H)$ by $U(\xi)(gH) := \xi(Hg^{-1})$ for all $\xi \in \ell^2(H\backslash G)$ and $g \in G$. It is clear that U is a unitary operator from $\ell^2(H\backslash G)$ onto $\ell^2(G/H)$. Using the fact that the map $\gamma \mapsto \gamma^{-1}$ is a bijection between the set $< H\backslash G >$ of right cosets onto the set $< G/H >$ of left cosets, one easily checks $U(\lambda_l(f)(\xi)) = \lambda_r(f)(U(\xi))$ for all $f \in \mathcal{H}(G, H)$ and $\xi \in \ell^2(H\backslash G)$. In other words, these actions are unitarily equivalent, and so they define isomorphic concrete C^*-algebras acting on two different Hilbert spaces.

We would like to use this opportunity to correct a typo in the text of [1] as well. In Line 4 of Page 183 of [1] the summation should be over G (instead of $< H\backslash G >$).

References

[1] **V. Shirbisheh,** Property (RD) for Hecke pairs. Mathematical Physics, Analysis and Geometry, vol. **15**, no. 2, (2012), 173–192.

E-mail address: shirbisheh@gmail.com